

DAY-SCALE VARIABILITY OF 3C 279 AND SEARCHES FOR CORRELATIONS IN GAMMA-RAY, X-RAY, AND OPTICAL BANDS

R. C. HARTMAN,^{1,2} M. VILLATA,³ T. J. BALONEK,⁴ D. L. BERTSCH,¹ H. BOCK,⁵ M. BÖTTCHER,^{6,7} M. T. CARINI,⁸ W. COLLMAR,⁹ G. DE FRANCESCO,³ E. C. FERRARA,¹⁰ J. HEIDT,⁵ G. KANBACH,⁹ S. KATAJAINEN,¹¹ M. KOSKIMIES,¹¹ O. M. KURTANIDZE,¹² L. LANTERI,³ A. LAWSON,¹³ Y. C. LIN,¹⁴ A. P. MARSCHER,¹⁵ J. P. MCFARLAND,¹⁰ I. M. MCHARDY,¹³ H. R. MILLER,¹⁰ M. NIKOLASHVILI,¹⁶ K. NILSSON,¹¹ J. C. NOBLE,¹⁵ G. NUCCIARELLI,¹⁷ L. OSTORERO,³ T. PURSIMO,¹¹ C. M. RAITERI,³ R. REKOLA,¹¹ T. SAVOLAINEN,¹¹ A. SILLANPÄÄ,¹¹ A. SMALE,¹⁸ G. SOBRITO,³ L. O. TAKALO,¹¹ D. J. THOMPSON,¹ G. TOSTI,¹⁷ S. J. WAGNER,⁵ AND J. W. WILSON¹⁰

Received 2001 March 1; accepted 2001 May 4

ABSTRACT

Light curves of 3C 279 are presented in optical (*R* band), X-rays (*RXTE*/*PCA*), and γ rays (*CGRO*/*EGRET*) for 1999 January–February and 2000 January–March. During both of those epochs the γ -ray levels were high and all three observed bands demonstrated substantial variation, on timescales as short as 1 day. Correlation analyses provided no consistent pattern, although a rather significant optical/ γ -ray correlation was seen in 1999, with a γ -ray lag of ~ 2.5 days, and there are other suggestions of correlations in the light curves. For comparison, correlation analysis is also presented for the γ -ray and X-ray light curves during the large γ ray flare in 1996 February and the two γ -bright weeks leading up to it; the correlation at that time was strong, with a γ -ray/X-ray offset of no more than 1 day.

Subject headings: gamma rays: observations — quasars: individual (3C 279)

1. INTRODUCTION

The discovery by the *EGRET* instrument on the *Compton Gamma Ray Observatory* (*CGRO*) that blazars can be strong γ -ray emitters posed an intriguing question: what is the mechanism responsible for this previously unknown and sometimes dominant high-energy emission? Obvious candidates are the well-known synchrotron self-Compton (SSC) and external-Compton (EC) models; in both of these suggested processes, the synchrotron-emitting relativistic electrons would energize soft photons via the inverse-Compton process. In this scenario, the principal matter of debate is the origin of the soft photons, the choices being synchrotron photons alone (SSC; e.g., Marscher & Gear 1985;

Maraschi, Ghisellini, & Celotti 1992; Bloom & Marscher 1996), photons of different provenance, i.e., accretion disk (ECD; Dermer, Schlickeiser, & Mastichiadis 1992; Dermer & Schlickeiser 1993; Sikora, Begelman, & Rees 1994) or broad-line region (ECR; Blandford & Levinson 1995; Ghisellini & Madau 1996; Dermer, Sturmer, & Schlickeiser 1997), or a combination of those possibilities.

Other possibilities, which have not yet been as carefully explored, are the proton-driven models. In the proton-initiated cascade (PIC) scenario (Mannheim 1993), very high-energy protons initiate photopion production, resulting in a γ -ray/electron/positron cascade. More recently, other proton-driven models have been discussed (e.g., Protheroe 1996a, 1996b; Rachen 2000; Aharonian 2000; Mücke & Protheroe 2001). These seem to be more readily applied to the lower-luminosity blazars (often described as high-frequency-peaked blazars, HBLs), but might also be adaptable for higher-luminosity blazars such as 3C 279.

Since the models predict different relationships between variations in the different observing bands, intensive simultaneous monitoring in several widely-spaced bands can provide crucial information on the radiation mechanisms and the structure of the jet. For that reason, several coordinated multiwavelength campaigns have been carried out in the last years on *EGRET*-detected blazars.

We present here the results of simultaneous monitoring in three bands, GeV γ -rays (from the *CGRO*/*EGRET* instrument), X-rays (from the *PCA* instrument on the *RXTE* satellite), and *R*-band optical (from a number of observers and ground-based observatories). For comparison, we also show a similar analysis of the γ -ray and X-ray light curves from the three weeks leading up to and including the large γ -ray flare in early February of 1996 (Wehrle et al. 1998). There was little optical coverage at that time, but the γ -rays and X-rays were strongly correlated.

2. OBSERVATIONS

The campaigns were centered around the following observation sequences:

¹ Code 661, NASA/GSFC, Greenbelt, MD 20771.
² rch@egret.gsfc.nasa.gov.
³ Osservatorio Astronomico di Torino, Strada Osservatorio 20, I-10025 Pino Torinese, Italy.
⁴ Department of Physics and Astronomy, Colgate University, 13 Oak Drive, Hamilton, NY 13346-1398.
⁵ Landessternwarte Königstuhl, 69117 Heidelberg, Germany.
⁶ Department of Space Physics and Astronomy, Rice University, Houston, TX 77005-1892.
⁷ Chandra Fellow.
⁸ Dept. of Physics and Astronomy, Western Kentucky University, 1 Big Red Way, Bowling Green, KY 42104.
⁹ Max-Planck-Institut für Extraterrestrische Physik, P.O. Box 1603, 85740 Garching, Germany.
¹⁰ Department of Physics and Astronomy, Georgia State University, Atlanta, GA 30303.
¹¹ Tuorla Observatory, Väisälantie 20, FIN-21500 Piikkiö, Finland.
¹² Astrophysikalisches Institut Potsdam, An der Sternwarte 16, 14482 Potsdam, Germany.
¹³ Department of Physics and Astronomy, University of Southampton, UK.
¹⁴ W. W. Hansen Experimental Physics Laboratory, Stanford University, Stanford, CA 94305.
¹⁵ Institute for Astrophysical Research, Boston University, 725 Commonwealth Ave., Boston, MA 02215.
¹⁶ Abastumani Observatory, 383762 Abastumani, Republic of Georgia.
¹⁷ Osservatorio Astronomico di Perugia, Via Bonfigli, 06123 Perugia, Italy.
¹⁸ Code 662, NASA Goddard Space Flight Center, Greenbelt, MD 20771.

1. 1996 January 16–February 6 (*CGRO* viewing periods 511.0, 511.5)
2. 1999 January 20–February 1 (*CGRO* viewing periods 806.5, 806.7)
3. 2000 February 9 – March 1 (*CGRO* viewing periods 910.0, 911.1)

CGRO viewing period 806.5 was an observation of 3C 273 during which EGRET was scheduled to be off. Because of the optical brightness of 3C 279, EGRET was turned on in full-field-of-view-mode to see if 3C 279 was detectable in the EGRET energy range. It was indeed bright, which led to the implementation of a target-of-opportunity observation, viewing period 806.7, with 3C 279 better centered in the field of view.

CGRO viewing period 910.0 was a 3C 279 target of opportunity, based on optical activity and brightness. Viewing period 911.1 was an observation of 3C 273 during which EGRET was scheduled to be off; due to the high γ -ray level seen in viewing period 910.0, EGRET was left on for viewing period 911.1 and was switched to full-field-mode to maximize the sensitivity to 3C 279, which was well off-axis.

2.1. Optical

R-band observations were made at a number of observatories, as described below, during both the 1999 and the 2000 campaigns. The coverage thus provided was the best ever obtained on a flat-spectrum radio quasar (FSRQ) during an EGRET observation. Most of the observations were made from European observatories. Several of the authors and observatories are members of the WEBT consortium.

3C 279 was observed during 1999 January 18–February 13 and 2000 January 29–February 19 at the Abastumani Astrophysical Observatory (Republic of Georgia) using a Peltier-cooled ST-6 CCD camera attached to the Newtonian focus of the 70 cm meniscus telescope (1/3). The full frame field of view is 14.9×10.7 arcmin². All observations are performed using combined filters of glasses which match well the standard *B*, *V* (Johnson) and *R_C*, *I_C* (Cousins) bands. Because the scale of the CCD and the meniscus telescope resolution are 2.3×2.7 arcsec² pixel⁻¹ and 1".5, respectively, the images are undersampled; therefore, the frames were slightly defocused to satisfy the sampling theorem. A full description of the Abastumani blazar monitoring program is given in Kurtanidze & Nikolashvili (1999).

Observations in 2000 were made with the 60 cm KVA telescope on La Palma, Canary Islands, using a ST-8 CCD camera with *BVR* filters. The data reduction was done using IRAF (with bias and flat-field corrections).

Observations were taken with the 1.2 m telescope of Calar Alto Observatory, Spain, and with the 0.7 m telescope of the Landessternwarte Heidelberg. Both telescopes are equipped with *LN₂*-cooled CCD cameras. Observations in Heidelberg are carried out with a Johnson *R*-band filter. The Calar Alto observations were carried out in Johnson *R* (in 2000) and R_{öser *R* (in earlier years). Standard debiasing and flat-fielding were carried out before performing differential aperture photometry.¹⁹}

¹⁹ Finding charts and comparison sequences are available at <http://www.lsw.uni-heidelberg.de/projects/extragalactic/charts.html> for 3C 279, along with many other blazars.

Observations were performed using Lowell Observatory's 42 inch Hall telescope and the 24 inch telescope of the Mount Stromlo/Siding Spring Observatories. Both telescopes are equipped with a direct CCD camera and an autoguider. The observations were made through *VRI* filters. Repeated exposures of 90 s were obtained for the star field containing 3C 279 and several comparison stars (Smith et al. 1985). These comparison stars were internally calibrated and are located on the same CCD frame as 3C 279. They were used as the reference standard stars in the data reduction process. The observations were reduced following Noble et al. (1997), using the method of Howell & Jacoby (1986). Each exposure is processed through an aperture photometry routine which reduces the data as if it were produced by a multistar photometer. Differential magnitudes can then be computed for any pair of stars on the frame. Thus, simultaneous observations of 3C 279, several comparison stars, and the sky background will allow one to remove variations which may be due to fluctuations in either atmospheric transparency or extinction. The aperture photometry routine used for these observations is the "phot" task in IRAF.

Observations were taken with the 2.5 m Nordic Optical Telescope (NOT) on La Palma, Canary Islands, Spain, using the ALFOSC instrument with a 2000 \times 2000 CCD camera (0".189 pixel⁻¹), and *V*- and *R*-filters. Data reduction (including bias and flat-field corrections) was made either with standard IRAF or MIDAS (J. Heidt) routines.

Observations at the Perugia Observatory were carried out with the Automatic Imaging Telescope (AIT). The AIT is based on an equatorially mounted 40 cm *f*/5 Newtonian reflector. A CCD camera and Johnson-Cousins *BVR_CI_C* filters are utilized for photometry (Tosti, Pascolini, & Fiorucci 1996). The data were reduced using aperture photometry with the procedure described in that reference.

Observations at the Torino Observatory were done with the 1.05 m REOSC telescope. The equipment includes an EEV CCD camera (1296 \times 1152 pixels, 0".467 pixel⁻¹) and standard (Johnson-Cousins) *BVRI* filters. Frames are reduced by the Robin procedure locally developed (Lanteri 1999), which includes bias subtraction, flat-fielding, and circular Gaussian fit after background subtraction. The magnitude calibration was performed according to the photometric sequence by Raiteri et al. (1998). Magnitudes were converted to fluxes by using a *B*-band Galactic extinction of 0.06 mag and following Rieke & Lebofsky (1985) and Cardelli, Clayton, & Mathis (1989).

2.2. X-Rays

3C 279 was the target for a series of 36 *RXTE* monitoring observations during 1999 January 2–February 16, for a total on-source time of 67 ks. The X-ray data presented here were obtained using the Proportional Counter Array (PCA) instrument in the Standard 2 and Good Xenon configurations, with time resolutions of 16 s and less than 1 μ s, respectively. Only PCUs 0, 1, and 2 were reliably on throughout the observations, and we limit our analysis to data from these detectors.

A further sequence of 28 monitoring observations was performed with *RXTE* in 2000 February, using the same instrumental configurations, for a total on-source time of 104 ks. For this sequence we utilized data from PCUs 0 and 2.




FIG. 1.—3C 279 light curves for early 1999 in γ -rays, X-rays, and R-band optical.

Data analysis was performed using *RXTE* standard analysis software, FTOOLS 5.0. Background subtraction of the PCA data was performed utilizing the “L7-240” models generated by the *RXTE* PCA team. The quality of the background subtraction was checked in two ways: (1) by comparing the source and background spectra and light curves at high energies (50–100 keV) where the source itself no longer contributes detectable events; and (2) by using the same models to background-subtract the data obtained during slews to and from the source.

2.3. Gamma Rays

The EGRET instrument is sensitive to γ -rays in the energy range 30 to 30,000 MeV. Its capabilities and calibration are described in Thompson et al. (1993); Esposito et al. (1999); and Bertsch et al. (2001). Point source data are analyzed using likelihood techniques (Mattox et al. 1996). The choice of 1 day as the unit of integration is dictated by the sensitivity of the EGRET detector and the general level of the emission during the time intervals of these observations. In TeV γ -rays, significant variations have been seen




FIG. 2.—3C 279 light curves for early 2000 in γ -rays, X-rays, and R-band optical.

on timescales well under 1 hr and may very well be present in 3C 279 in GeV γ -rays also.

The 1999 and 2000 γ -ray data presented here have been shown previously in a preliminary form in Hartman et al. (2001a, 2001b). Unfortunately, in both of those references, there was a 1 day error in the Julian Dates for the γ -ray observations. Thus, the ~ 3.5 day optical to γ -ray lag in 1999 tentatively reported there corresponds to the ~ 2.5 day lag discussed here.

2.4. Light Curves

The light curves resulting from the observations described above are shown in Figures 1 and 2.

3. CORRELATION ANALYSIS AND RESULTS

Although the γ -ray observations were continuous, both the R band and X-rays were only sampled, sometimes irregularly and/or sparsely; thus, the discrete correlation function (DCF), which was designed for analysis of unevenly sampled data (Edelson & Krolik 1988), was used for this analysis. Because of the large statistical errors on the




FIG. 3.—3C 279 correlation functions (DCF) for 1999

EGRET data points, the initial analysis was done using equation (3) of Edelson & Krolik (1988); this resulted in unphysical normalization in the correlation results. According to J. Krolik (2000, private communication), this is a known but unresolved effect in DCF analyses. Reanalysis ignoring the errors on the EGRET data points resulted in reasonable normalizations and produced equally significant correlations. Therefore, the correlation results shown below all ignore the EGRET errors.

Figures 3 and 4 show the results of the DCF analyses for the 1999 and 2000 light curves, respectively. The following is a summary of the evidence found for correlations.

3.1. 1999

3.1.1. *Gamma Ray/X-Ray*

Two possible correlations are found, in which the γ -rays lag the X-rays by about 10 and 5–6 days. The first links the highest two X-ray points with the two γ -ray peaks; the second links the first γ -ray peak with the second X-ray high point and the second (sharp) γ -ray peak with a time period with little X-ray coverage. While mathematically possible, these correlations are unconvincing because of the very limited X-ray coverage around the relevant times. In addition, the long delays are probably difficult to account for theoretically.




FIG. 4.—3C 279 correlation functions (DCF) for 2000

3.1.2. *Gamma Ray/Optical*

If a 2–3 day γ -ray lag is assumed, the γ -ray light curve is very similar to that in the R band, and the DCF analysis finds this to be a rather strong correlation. It is the most convincing correlation found in the six DCF analyses for 1999 and 2000.

A negative correlation with an 8–9 day optical lag links the first γ -ray peak with the R-band minimum around TJD 210 and the second γ -ray peak in the optically unsampled TJD 213–218 interval. In addition to being unconvincing because of coverage limitations, this seems quite unphysical.

3.1.3. *X-Ray/Optical*

A possible correlation with a 2.5 day optical lag requires that the two optical peaks around TJD 194 and 198 be linked to the two highest X-ray points and ignores the strongest and sharpest optical feature, placing it in the weak X-ray minimum of TJD 202. This is unconvincing because of the very limited coverage around the two X-ray high points and also because the most prominent optical feature is ignored.

A fairly significant correlation with a 7–8 day optical lag links the two highest X-ray points with the optical peaks at about TJD 198.5 and 204.0. This is conceivable, but in addition to the poor X-ray sampling around the relevant times, the 7–8 day offset seems difficult to accommodate theoretically.




FIG. 5.—3C 279 light curves for early 1996 in γ -rays and X-rays

3.2. 2000

3.2.1. Gamma Ray/X-Ray

At zero time-delay, there is a sharp peak in the DCF; its statistical significance is only about 1.8σ , but the correlated pattern is obvious to the eye in the light curves, not only around the sharp peaks, but in the entirety of both light curves.

The variations seen here are more complicated than those seen in both γ -rays and X-rays in 1996 January–February, when a zero time-delay was also seen.

3.2.2. Gamma Ray/Optical

No significant correlation was found.

3.2.3. X-Ray/Optical

A possible correlation with a 1 day optical lag seems plausible, but ignores the X-ray peak around TJD 585.

Another possible correlation is seen with an 11 day X-ray lag. This is rather unconvincing because of X-ray variations that do not show up in the optical and is also somewhat implausible physically.




FIG. 6.—3C 279 γ -ray/X-ray correlation function (DCF) for 1996

3.3. 1996 High State and Large Flare

For comparison, DCF analysis is presented of the γ -ray and X-ray light curves for the three weeks in 1996 January–February leading up to and including the large γ -ray flare. The light curves, adapted from Wehrle et al. 1998, are shown in Figure 5. To the eye, the γ -rays and X-rays appear well-correlated, with no apparent lag; this is confirmed by the DCF analysis, the results of which are shown in Figure 6.

4. DISCUSSION AND CONCLUSIONS

In the 1999 and 2000 light curves, the correlation that is most apparent to the eye is that in 1999 between the γ -rays and the optical, with a ~ 2.5 day γ -ray lag. With the γ -rays, peaked around TJD 206.5, apparently correlating with the optical peak around TJD 204, it is not easy to imagine a scenario that could produce such a sequence (a discussion on various possibilities can be found in Hartman et al. 2001a, 2001b).

Another correlation that seems apparent to the eye, that of the X-rays and γ -rays in 2000, gives a disappointingly weak effect ($\sim 2 \sigma$ for zero delay) when analyzed with the DCF, as noted above. Examination of the autocorrelations in those two bands (Fig. 7) provides some assistance in interpreting the results. Although the X-ray autocorrelation is fairly routine, that for the γ -rays is unusual. Apparently this is due to the two 1 day high points separated by a day for which the best γ -ray flux estimate is zero, albeit with substantial statistical errors on all of the points. (Examination of the photon maps for the 3 days under discussion verifies that the γ -rays do disappear during the middle day.)

Thus, there is no consistent pattern found. This could be because the emission in the three bands investigated really does have no persistent relationship or merely because the




FIG. 7.—3C 279 γ -ray and X-ray autocorrelations for 2000

data are not adequate, in coverage and/or statistical accuracy, to bring out such relationships.

What correlations and time delays are to be expected here? Detailed predictions about the theoretically expected light curves are difficult because of the multitude of physical processes potentially involved in the formation and evolution of the particle and photon spectra, in particular in FSRQs. While detailed modeling of variability patterns expected in high-frequency peaked BL Lac objects (which are well modeled with pure SSC models) has been done (e.g., Takahashi et al. 1996; Georganopoulos & Marscher 1998; Kusunose, Takahara & Li 2000; Li & Kusunose 2000), detailed theoretical work relevant to the short-term variability of the multicomponent spectra probably present in the high-energy emission from FSRQs is still in its very early stages (see, e.g., Sikora et al. 2001). Thus, we must restrict the discussion of the expected time lags to order-of-magnitude estimates at this point.

Frequency-dependent time lags in the short-term variability of FSRQs like 3C 279 are likely to be related to either the electron cooling in the effectively emitting region or the dynamical timescale on which the soft seed photon fields for Compton scattering are changing in the frame of the relativistically moving emitting region. The timescale for acceleration of relativistic electrons might be of the order of the timescale for Fermi acceleration, $\tau_{\text{acc}} \sim 2\pi r_L/c \sim 3.6 \times 10^{-7} \gamma B$ s in the comoving frame of the emitting region (where r_L is the Larmor radius, γ is the electron Lorentz factor, and B is the magnetic field in G) or $\tau_{\text{acc}}^* = \tau_{\text{acc}}/D$ in the observer's frame (where $D \sim 10$ is the Doppler boosting factor determining the time contraction between the comoving and the observer's frame). Thus, for any reasonable value of the magnetic field, variability on the acceleration timescale will be smeared out by light travel time effects and would be too short to be resolvable with current multi-wavelength observations anyway.

Based on the multiepoch multiwavelength spectral fits to 3C 279 presented in Hartman et al. (2001c), we can estimate the typical electron cooling timescale in the emitting region, assuming that (as indicated by the spectral fits) electron cooling is dominated by inverse-Compton scattering of external radiation fields. Taking into account both the contributions from direct accretion disk radiation and from reprocessing of this radiation within the broad line region, we find the observed cooling time is

$$\tau_{\text{cool}}^* = \left(\frac{4}{3} c \sigma_T \frac{u_s}{m_e c^2} \gamma D \right)^{-1},$$

where σ_T is the Thomson cross section and the energy density of soft photons in the emitting region is given by

$$u_s \approx \frac{L_D}{4\pi c} \left(\frac{1}{z^2 \Gamma^2} + \frac{\Gamma^2 \tau_{\text{blr}}}{r_{\text{blr}}^2} \right).$$

Here L_D is the accretion disk luminosity, z is the distance of the emitting region from the central engine, Γ is the bulk Lorentz factor of the emitting region, and τ_{blr} and r_{blr} are the radial Thomson depth of the broad line region and its average distance from the central engine, respectively. For reasonable values of the parameters ($L_D \sim 10^{46}$ ergs s $^{-1}$, $z \sim 0.025$ pc, $\Gamma \sim 10$, $\tau_{\text{blr}} \sim 0.003$, and $r_{\text{blr}} \sim 0.1$ pc), we find that the cooling timescale relevant to electrons emitting in the EGRET energy regime ($\gamma \gtrsim 10^4$) is of the order of 1 to

several hours, while for particles emitting predominantly at X-ray and optical frequencies ($\gamma \lesssim 100$), it is expected to be 1 to several days, which would then be the relevant timescale determining time lags between different energy bands.

If time lags are dominated by the dynamical timescale on which the soft seed photon fields (for Compton scattering, to produce the high-energy radiation) are changing, we would expect typical time lags of

$$\tau_{\text{dyn}}^* \sim \frac{\Delta z(1 - \beta_r \cos \theta)}{c}.$$

For typical values of the parameters ($\Delta z \sim 0.1$ pc, $D \sim \Gamma \sim 10$, and $\cos \theta \sim \beta_r$), this gives delays of a few days. Thus, in both cases the expected time lags between different photon energy bands would be of the order of 1 to a few days.

In HBLs, proton-driven models can produce time delays of the order of a day or less (A. Mücke 2001, private communication). This is probably true also of the more complicated FSRQs such as 3C 279, but so far there has been no theoretical study demonstrating this.

It should be noted that, despite the similar flux levels and variations in 1999 and 2000, Hartman et al. (2001c) have shown evidence that conditions in the inner region of 3C 279 may have been substantially different in 2000 than in 1999. In particular, the strength of the broad line emission may have been much weaker in 2000.

Several strong implications for future investigations such as this are clear:

γ -rays.—Two necessary improvements are obvious: considerably better statistics and longer observation intervals. Both of these requirements will be met by the *GLAST* mission, planned for launch in 2006: (1) The *GLAST* observation plan (at least for the first year or two) is to operate in a scanning mode, so that a large fraction of the sky will be covered during each orbit. Objects near the equator, such as 3C 279, will receive good exposure on every orbit during the scanning part of the mission, and the entire sky will receive significant coverage over 1 day. (2) Its larger effective area, larger field of view, and better point-spread function (compared with EGRET) will provide much better statistics and sensitivity than are available from EGRET.

The Italian γ -ray telescope *AGILE* (Vercellone et al. 1999), planned for launch in 2002–2003, will have sensitivity comparable to that of EGRET, but with significantly better angular resolution. Its observing program will permit longer observations than were usually possible with EGRET.

X-rays.—The obvious need here is for more uniform coverage. Unfortunately, it is unlikely that the *RXTE* satellite, with its very flexible scheduling, will be operating by the time of the *GLAST* launch. The big X-ray missions expected to be operating in parallel with *GLAST* are likely to be less flexible and accessible than *RXTE*. They will, however, have much greater sensitivity, permitting investigation of dimmer objects than *RXTE*. The X-ray monitors on *HETE II* and *Swift* may be able to provide some assistance, but their availability and applicability are unclear at the present time. Thus, good X-ray time-sampling during the *GLAST* era appears uncertain at present.

Optical.—Although the optical coverage was good most of the time during and around the EGRET observations used here, there were some substantial holes that allowed

the DCF to suggest unlikely correlations. Future investigations of this type will certainly need to improve upon this. Some of the intensive optical monitoring presented here utilized automated telescopes. Hopefully, additional automated systems, at sites throughout the world, will be available by the time of the *GLAST* launch.

The WEBT (Whole Earth Blazar Telescope; Mattox 1999a, 1999b; Villata et al. 2000) is a different approach to intensive optical monitoring. It is a consortium of about 20 optical observatories around the world, formed to facilitate 24 hour high-time-density blazar observations during multiwavelength campaigns.²⁰

The work at Torino Observatory and Perugia University Observatory was partly supported by the Italian Ministry for University and Research (MURST) under grant Cofin98-02-32 and by the Italian Space Agency (ASI).

The Nordic Optical Telescope is operated on the island of La Palma jointly by Denmark, Finland, Iceland, Norway, and Sweden in the Spanish Observatorio del Roque de Los Muchachos of the Instituto de Astrofísica de

Canarias. The Tuorla Observatory authors wish to thank The Finnish Academy for support.

H. B., J. H., and S. J. W. acknowledge support by the DFG (SFB 328 and 439) and CAHA/DSAZ for support during several observing runs on Calar Alto.

O. M. K. thanks the Astrophysikalisches Institute Potsdam for support.

The Georgia State University authors wish to thank Lowell and Mount Stromlo/Siding Spring Observatories for allocations of observing time. This work has been supported in part by an award from GSU's RPE Fund to PEGA and by grants from the Research Corporation and NASA (NAGW-4397).

The work of M. B. is supported by NASA through Chandra Postdoctoral Fellowship Award No. 9-10007, issued by the Chandra X-ray Center, which is operated by the Smithsonian Astrophysical Observatory for and on behalf of NASA under contract NAS 8-39073.

IRAF is distributed by the National Optical Astronomy Observatories, which is operated by the Association of Universities for Research in Astronomy, Inc., under cooperative agreement with the National Science Foundation.

²⁰ For additional information see <http://astro.fmarion.edu/webt>.

REFERENCES

- Aharonian, F. A. 2000, *NewA*, 5, 377A
 Bertsch, D. L., Hartman, R. C., Hunter, S. D., Thompson, D. J., & Sreekumar, P. 2001, in *Gamma 2001: Gamma-Ray Astrophysics 2001*, ed. S. Ritz, N. Gehrels, & C. Schrader (New York: AIP), in press
 Blandford, R. D., & Levinson, A. 1995, *ApJ*, 441, 79
 Bloom, S. D., & Marscher, A. P. 1996, *ApJ*, 461, 657
 Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, *ApJ*, 345, 245
 Dermer, C. D., & Schlickeiser, R. 1993, *ApJ*, 416, 458
 Dermer, C. D., Schlickeiser, R., & Mastichiadis, A. 1992, *A&A*, 256, L27
 Dermer, C. D., Sturmer, S. J., & Schlickeiser, R. 1997, *ApJS*, 109, 103
 Edelson, R. A., & Krolik, J. H. 1988, *ApJ*, 333, 646
 Esposito, J. A., et al. 1999, *ApJS*, 123, 203
 Georganopoulos, M., & Marscher, A. P. 1998, *ApJ*, 506, L11
 Ghisellini, G., & Madau, P. 1996, *MNRAS*, 280, 67
 Hartman, R. C., et al. 2001a, *Mem. Soc. Astron. Ital.*, in press
 Hartman, R. C., et al. 2001b, in *ASP Conf. Ser. 224, Probing the Physics of Active Galactic Nuclei by Multiwavelength Monitoring*, ed. B. M. Peterson, R. S. Polidan, & R. W. Pogge (San Francisco: ASP), 249
 Hartman, R. C., et al. 2001c, *ApJ*, 553, 683
 Howell, S. B., & Jacoby, G. J. 1986, *PASP*, 98, 802
 Kurtanidze, O. M., & Nikolashvili, M. G. 1999, in *Proc. OJ-94 Annual Meeting 1999, Blazar Monitoring towards the Third Millennium*, ed. C. M. Raiteri, M. Villata, & L. O. Takalo (Pino Torinese: Osservatorio Astronomico di Torino), 25
 Kusunose, M., Takahara, F., & Li, H. 2000, *ApJ*, 536, 299
 Lanteri, L., 1999, in *Proc. OJ-94 Annual Meeting 1999, Blazar Monitoring towards the Third Millennium*, ed. C. M. Raiteri, M. Villata, & L. O. Takalo (Pino Torinese: Osservatorio Astronomico di Torino), 125
 Li, H., & Kusunose, M. 2000, *ApJ*, 536, L729
 Mannheim, K. 1993, *A&A*, 269, 67
 Maraschi, L., Ghisellini, G., & Celotti, A. 1992, *ApJ*, 397, L5
 Marscher, A. P., & Gear, W. 1985, *ApJ*, 298, 114
 Mattox, J. R., et al. 1996, *ApJ*, 461, 396
 Mattox, J. R., et al. 1999a, in *Proc. OJ-94 Annual Meeting 1999, Blazar Monitoring towards the Third Millennium*, ed. C. M. Raiteri, M. Villata, & L. O. Takalo (Pino Torinese: Osservatorio Astronomico di Torino), 44
 Mattox, J. R., et al. 1999b, in *ASP Conf. Ser. 189, CCD Precision Photometry Workshop*, ed. E. R. Craine, R. A. Tucker, & J. Barnes (San Francisco: ASP), 95
 Mücke, A., & Protheroe, R. J. 2001, *Astropart. Phys.*, 15, 121
 Noble, J. C., Carini, M. T., Miller, H. R., & Goodrich, B. 1997, *AJ*, 113, 1995
 Protheroe, R. J. 1996a, *Adelaide Univ. preprint ADP-AT-96-4*
 ———. 1996b, *Adelaide Univ. preprint ADP-AT-96-7*
 Rachen, J. P. 2000, in *AIP Conf. Proc. 515, GeV-TeV Astrophysics: Toward a Major Atmospheric Cherenkov Telescope V*, ed. B. L. Dingus, M. H. Salamon, & D. B. Kieda (New York: AIP), 41
 Raiteri, C. M., Villata, M., Lanteri, L., Cavallone, M., Sobrito, G. 1998, *A&AS*, 130, 495
 Rieke, G. H., & Lebofsky, M. J. 1985, *ApJ*, 288, 618
 Sikora, M., Begelman, M. C., & Rees, M. J. 1994, *ApJ*, 421, 153
 Sikora, M., Blazewski, M., Begelman, M. C., & Moderski, R., 2001, *ApJ*, in press
 Smith, P. S., Balonek, T. J., Heckert, P. A., Elston, R., & Schmidt, G. D. 1985, *AJ*, 90, 1184
 Takahashi, T., et al. 1996, *ApJ*, 470, L89
 Thompson, D. J., et al. 1993, *ApJS*, 86, 629
 Tosti, G., Pascolini, S., & Fiorucci, M. 1996, *PASP*, 108, 706
 Vercellone, S., et al. 1999, in *Proc. OJ-94 Annual Meeting 1999, Blazar Monitoring towards the Third Millennium*, ed. C. M. Raiteri, M. Villata, & L. O. Takalo (Pino Torinese: Osservatorio Astronomico di Torino), 138
 Villata, M., et al. 2000, *A&A*, 363, 108
 Wehrle, A., et al. 1998, *ApJ*, 497, 178