


# ANABOLISM II:

## Biosynthesis of Fatty Acids & Lipids

### ANABOLISM II: Biosynthesis of Fatty Acids & Lipids



# Fatty Acid Biosynthesis

Catabolism  
Anabolism


- Contrast with Sugars
  - Lipids have hydro-carbons not carbo-hydrates
  - more reduced=more energy
  - Long-term storage vs short-term storage
  - Lipids are essential for structure in ALL organisms: membrane phospholipids
- Catabolism of fatty acids
  - produces acetyl-CoA
  - produces reducing power (NADH, FADH<sub>2</sub>)
  - takes place in the mitochondria
- Anabolism of fatty acids
  - requires acetyl-CoA and sufficient carbohydrates
  - requires reducing power from NADPH
  - takes place in cytosol in animals, chloroplast in plants

Anabolism of fat will occur only when there is excess, carbon, electrons, and  $\Delta G$

Acetyl-CoA & carbohydrates → NADPH → ATP

How is this monitored in the cell?

# Fatty Acid Biosynthesis



Both of these compartments are where there are ample Sources of NADPH:

in adipocytes: pentose phosphate pathway and malic enzyme

NADPH is made as malate converts to pyruvate + CO<sub>2</sub>.

in hepatocytes and mammary gland: pentose phosphate pathway

NADPH is made as glucose-6-phosphate converts to ribulose 6-phosphate.

in plants: photosynthesis

# Fatty Acid Biosynthesis


## EXAMPLE: Synthesis of Palmitate (16:0)

Where do the carbons come from? **Acetyl-CoA**

1. 8 acetyl-CoA x 2 carbons = 16 carbons (palmityl-CoA)
2. Longer fatty acids and desaturases use palmityl-CoA
3. 1 acetyl-CoA "primes" the enzyme
  - a. other "acetyl-CoA-derived units" are ACTIVATED by carboxylation (recall gluconeogenesis)
  - b. Used to make 7 malonyl-CoAs... using? **ATP & CO<sub>2</sub>**




2. Seven cycles of condensation, reduction, dehydration, and reduction... using 2 **NADPH** to reduce the  $\beta$ -keto group and trans-double bond



How is the acetyl-CoA transported out of mitochondria?



# Fatty Acid Biosynthesis



# Fatty Acid Biosynthesis

OMSGAP

- Fatty acids are built in several passes, processing **one acetate unit** at a time.
- The acetate is coming from activated malonate in the form of **malonyl-CoA**.
- Each pass involves reduction of a **carbonyl** carbon to a **methylene** carbon.




here is the third ATP!

- Making the **Malonyl-CoA**:
  - Reaction carboxylates acetyl-CoA
  - Catalyzed by **acetyl-CoA carboxylase (ACC)**

## Fatty Acid Biosynthesis

### Acetyl-CoA Carboxylase Reaction:


- The enzyme has three subunits:
  - One unit has biotin covalently linked to Lys.
  - Another subunit is biotin carboxylase
  - The third subunit is a transcarboxylase
  - Biotin carries  $\text{CO}_2$ .
  - In animals, all three subunits are on one polypeptide chain.
- $\text{HCO}_3^-$  (bicarbonate) is the soluble source of  $\text{CO}_2$ .
- Two-step Rxn similar to carboxylations catalyzed by *pyruvate carboxylase* and *propionyl-CoA carboxylase*
  - $\text{CO}_2$  binds to biotin.
    - Reaction with ATP produces carboxy-phosphate.
    - Activated  $\text{CO}_2$  is attached to N in ring of biotin.
  - Enzyme undergoes conformational change to carry carbamoyl to transcarboxylase site
  - $\text{CO}_2$  attaches to acetyl-CoA, which leaves active site



# Fatty Acid Biosynthesis

## Fatty Acid Synthase (FAS)

- Catalyzes a repeating four-step sequence that elongates the fatty acyl chain by two carbons at each step
  - NADPH as the electron donor
  - **Condensation** with acetate
    - $\beta$ -ketoacyl-ACP synthase (KS)
  - **Reduction** of  $\beta$ -carbonyl to hydroxyl
    - $\beta$ -ketoacyl-ACP reductase (KR)
  - **Dehydration** of alcohol to alkene
    - $\beta$ -hydroxyacyl-ACP dehydratase (DH)
  - **Reduction** of alkene to alkane
    - enoyl-ACP reductase (ER)
  - **Chain transfer/charging**
    - malonyl/acetyl-CoA ACP transferase (M/AT)
- Overall goal: attach acetate unit (2-carbon) from malonyl-CoA to a growing chain and then reduce it.
- Reaction involves cycles of four enzyme-catalyzed steps:
- condensation of the growing chain with activated acetate (malonyl-CoA)
  - reduction of carbonyl to hydroxyl
  - dehydration of alcohol to trans-alkene
  - reduction of alkene to alkane
- The growing chain is initially attached to the enzyme via a thioester linkage on the acyl-carrier protein (ACP).
- What is this "ACP"?  
...let's look at the structure.



# Fatty Acid Biosynthesis

## FAS I vs. FAS II



### FAS I

- Single polypeptide chain in vertebrates
- Leads to single product: palmitate 16:0
- C-15 and C-16 are from the acetyl-CoA used to prime the Rxn
- FAS I in vertebrates and fungi

### FAS II

- Made of separate, diffusible enzymes
- Makes many products (saturated, unsaturated, branched, many lengths, etc.)
- Mostly in plants and bacteria

### FAS I




KS MAT DH ER KR ACP TE

What you can't see is the ACP.....

# Fatty Acid Biosynthesis


## Acyl Carrier Protein (ACP)



The *E. coli* ACP is a small 77-residue protein with a covalently attached prosthetic group, 4'-phosphopantetheine, at a Ser residue.

– In vertebrate FAS, it's a domain with a flexible arm to tether the growing acyl chain

- **Delivers acetate** (in the first step) or **malonate** (in all the next steps) to the fatty acid synthase enzymes
- **Shuttles the growing chain** from one active site to another during the four-step reaction



## Priming FAS

- Two thiols must be **charged with the correct acyl groups** before the condensation reaction can begin.
  - thiol from 4-phosphopantetheine in **ACP**
  - thiol from Cys  **$\beta$ -ketoacyl-ACP synthase (KS)**
- The acetyl group of acetyl-CoA is transferred to ACP.
  - catalyzed by **malonyl/acetyl-CoA transferase (MAT)**
  - **ACP** passes this acetate to the Cys of the **KS** domain of FAS 1.
  - **ACP** -SH group is recharged with **malonate** from **malonyl-CoA** again catalyzed by **MAT**

